


AFFIDAVIT

The undersigned Serena Gitto of Barzanò & Zanardo Roma S.p.A., 26 Via Piemonte, Rome, Italy hereby declares to be familiar with the English and Italian languages and that the attached is a true translation of the priority document.

A handwritten signature in black ink, appearing to read "Serena Gitto". The signature is fluid and cursive, with the first name "Serena" written in a larger, more prominent script than the last name "Gitto".

PURIFICATION, CLONING AND BIOCHEMICAL CHARACTERIZATION
OF XendoU, ENDORIBONUCLEASIC ACTIVITY INVOLVED IN SMALL
NUCLEAR RNA SPLICING-INDEPENDENT BIOSYNTHESIS IN
XENOPUS LAEVIS

5

Endoribonucleases play essential role in RNA metabolism participating both in "degradative" pathways, such as mRNA decay (Schoenberg and Chemokaiskaya, 1997), and in "maturative" pathways, to generate functional RNA molecules from primary transcripts. Only a few
10 endoribonucleases have been isolated in eukaryotes, most of them being involved in biosynthesis of translation apparatus components. Among these, there are ribonucleoprotein enzymes, such as RNase P and RNase MRP, which act as site-specific endoribonucleases: RNase P is involved in generation of the 5' end of tRNAs (Nashimoto, 1995), whereas RNase
15 MRP is implicated in processing of pre-rRNA (Lygerou et al., 1996). Other endonucleolytic activities, as 3'-tRNase, tRNA splicing endonuclease and members of RNase III family (Trotta et al., 1997; Bujnicki and Rychlewski, 2000; Zamore, 2001) are protein enzymes. 3'-tRNase is an eukaryotic spermidine-dependent endoribonuclease which precisely removes the 3'-
20 end tail from tRNA precursors (Castano et al., 1985; Nashimoto, 1995). tRNAs splicing endoribonucleases are required for the intron removal from pre-tRNAs: it is an Mg^{++} -dependent enzyme and cleaves pre-tRNAs at 5' and 3' splice sites, releasing products having 2'-3' cyclic phosphate and 5'OH ends (Peebles et al., 1983). RNases III are endoribonucleases acting
25 on double-strand RNA found in bacteria and eukaryotes: they were first isolated in *E. coli* (Court, 1993; Nicholson, 1997) and subsequently eukaryotic orthologs were identified on the basis of sequence similarity. *S. cerevisiae* RNase III (Rnt1p), was shown to be involved in several biosynthetic events such as pre-rRNAs, snRNAs and snoRNAs processing
30 (Elela et al., 1996; Kufel et al., 1999; Chanfreau et al., 1997; Allmang et al., 1999; Chanfreau et al., 1998). Recently it was shown that Rnt1p is also involved in the release of the intron-encoded snoRNAs, U18 and snR38 from their pre-mRNAs (Giorgi et al., 2001). Furthermore, a new member of eukaryotic RNases III family, named "Dicer", has been
35 identified in metazoa (Bemstein et al., 2001). It is known to be involved in interference (RNAi) pathway, generating 21-23 nt small interfering RNAs (siRNAs) from longer partially double stranded precursors. These processing products act in RNA-mediated gene regulation (Ambros,

2001). Cleavage by Rnase III releases 3'OH and 5' phosphate ends and it is Mg^{++} -dependent.

5 The authors of the invention previously demonstrated that an endoribonucleolytic activity plays a key role in the biosynthesis of the box C/D U16 snoRNA, encoded in the third intron of the L4 ribosomal protein gene in *X. laevis* (Caffarelli et al., 1994). The authors already showed that U16 processing from the host intron is alternative to the splicing reaction: thereby, synthesis of the L4 mRNA is alternative to the production of U16 snoRNA (Caffarelli et al., 1996). In this context the biosynthetic
10 mechanism of U16, per se, regulates the expression of L4 gene at the post-transcriptional level.

The authors now purified to homogeneity from *X. laevis* oocyte nuclear extracts (ONE) and characterized the endonucleolytic activity (XendoU, GenBank TM/EBI Data Bank AJ507315) responsible for the
15 processing of U16 snoRNA from its host intron. Partial protein sequencing allowed to clone XendoU cDNA, to express it and to study features of the enzyme. This protein represents a novel endoribonucleasic activity, being: i) poly-U specific, ii) single filament specific, iii) Mn^{++} -dependent and iv) able to release cleavage products with 5' OH and 2'-3' cyclic phosphate
20 ends.

It is therefore an object of the present invention a nucleic acid encoding for a protein with endoribonucleasic activity which is poliU sequences specific, Mn^{++} -dependent and releases as cleavage products
25 molecules having 2'-3' cyclic phosphate and 5'OH ends. Preferably nucleic acid includes substantially nucleotide sequence of figure 4, functional homologs thereof or complementary sequence thereto.

It is a further object of the invention a recombinant vector able to express effectively the inventive nucleic acid in prokaryotes.

30 It is a further object of the invention a recombinant vector able to express effectively the inventive nucleic acid in eukaryotes

Those skilled in the art will be able to recognize the most suitable vectors also considering the host organism in order to express the inventive nucleic acid.

35 It is a further object of the invention a protein with endoribonucleasic activity which is poly-U sequences specific, Mn^{++} -dependent and releases RNA molecules bearing 2',3' cyclic phosphate and 5'OH ends as cleavage products, or functional portions thereof.

Preferably the protein is encoded by the nucleic acid of the invention, more preferably protein substantially has amino acid sequence of figure 4. Advantageously the protein is produced by synthetic or recombinant route using methods known by those skilled in the art.

5 The invention will be now described without any limitation thereof referring to experimental procedures wherein reference will be made to the following figures:

Figure 1 shows activity assay and purification of XendoU. A, schematic representation of U16 snoRNA processing. P indicates U16-containing precursor (003 RNA); I-1a and I-1b represent the products generated by cleavage upstream from U16 while I-2a and I-2b represent their 3' complementary molecules thereto; 1-3 depicts the product originated from cleavage downstream from U16 while 1-4 represent its 3' respective complementary molecule; pre-U16 represents U16 precursor with additional 5' and 3' flanking sequences. Cap structure is shown as a black dot, exons as boxes, the intron as a continuous line and U16 snoRNA coding region as a thicker line. Large arrows localise the major sites of cleavage and small arrows the minor ones. B, *in vitro* U16 processing in unfractionated oocyte nuclear extracts (ONE) and with purified XendoU (XendoU). ³²P-labelled 003 RNA was incubated for the times indicated below; RNA was then extracted and separated by 6% polyacrilamide-7M urea gel. The specific cleavage products are indicated aside. C, scheme of procedure employed for XendoU purification from ONE. D, proteins from the active fractions during the purification were separated on SDS-PAGE and visualised by Blue Coomassie staining. Numerals below refer to the corresponding purification steps schematised in panel C. The arrow points to the purified enzyme with an apparent molecular mass of 37 kDa.

Figure 2 shows XendoU cleavage which requires Mn⁺⁺ cations. ³²P-labelled 003 RNA was incubated in oocyte nuclear extract (ONE) or with purified XendoU (XendoU) in the presence of different metal ions at a concentration of 6 mM. After 30 min, the reaction was stopped and the processing products analysed on 6% polyacrilamide-7M urea gel. In the lane - the RNA substrate was incubated with XendoU in the absence of ions. The specific cleavage products are schematised on the left side .

Figure 3 shows that the XendoU activity is U-specific and produces 2'-3' cyclic phosphate. A, ³²P ATP labelled synthetic

oligoribonucleotide P1, containing the distal cleavage site upstream from U16, and its mutant derivatives (P2, P3) were incubated with the unfractionated extract (lanes 2) or with purified XendoU (lanes 3), under standard conditions for 30 min. RNA molecules were extracted and analysed on 10% polyacrylamide-7M urea gel. In lanes 1 untreated RNA is shown, in lane M RNA marker generated by alkaline digestion of P1 is shown; arrows indicate cleavage sites. On the right side the sequences of the oligoribonucleotides are reported. B, The ^{32}P -labelled I-1 molecules, schematically represented on the left side, generated by incubation of U16-containing precursor in ONE (ONE), with purified XendoU (XendoU) or after injection in oocytes (*in vivo*), were gel purified, and their 3'end analysed. The molecules were incubated with 1 unit of alkaline phosphatase (lane 1) or with 10 mM HCl (lanes 2) or with alkaline phosphatase after acid treatment (lanes 3). After incubation the RNAs extracts were analysed by electrophoresis on 10 % polyacrylamide-7M urea gel. Untreated molecules were run as control in lanes 4.

Figure 4 shows cDNA and amino acid sequence of XendoU. Nucleotides of the 5' and 3' untranslated regions are shown in small letters, nucleotides of the ORF in capital letters. Above each codon the corresponding amino acid is shown. The sequence portions determined by automated Edman degradation and mass mapping experiments (see "Experimental Procedures") are indicated by numbers 1, 2 and 3. The stop codon is identified by an asterisk. Numbers on the right side of diagram correspond to the amino acid residues. Underlined are the amino acid sequences identified by MALDI-mapping experiments.

Figure 5 shows the functional analysis of *in vitro* translated XendoU A, SDS-PAGE analysis of [^{35}S] Methionine-labelled XendoU (lane 2) and of control luciferase (lane 1) produced by *in vitro* transcription and translation. The arrow points to XendoU protein. B, 003 RNA was incubated, in the presence of Mn^{++} ions, in ONE (ONE), with XendoU produced by reticulocyte lysate (ret-XendoU/+Mn), or with reticulocyte lysate as such (lane ret/+Mn). As control 003 RNA was incubated, in the absence of Mn^{++} ions, with XendoU produced by reticulocyte lysate (lanes ret-XendoU/-Mn). The numbers below indicate incubation times: 0 min (lane 1), 45 min (lanes 2).

Figure 6 shows that XendoU is involved in U86 snoRNA biosynthesis. A, U86 processing is analysed *in vivo* by injection of ^{32}P -

labelled U86-containing precursor (P) in *X. laevis* oocytes (lanes *in vivo*), or *in vitro* by incubation of the RNA precursor in ONE (ONE), or with purified XendoU (XendoU). The numbers below indicate different incubation times: 0 min (lane 1), 10 min (lanes 2), 45 min (lanes 3), 3 hours (lanes 4), 16 hours (lanes 5). RNA was then extracted and loaded on a 6% polyacrilamide-7M urea gel. The processing products are schematised aside. Arrows indicate specific XendoU cleavages. B, ³²P-labelled UhindIII primer, depicted below, was reacted with unlabelled 1-4 molecules obtained after 10 min of incubation in oocytes (lane *in vivo*), 45 min of incubation in ONE (ONE) or 45 min of incubation with purified XendoU (XendoU). The products of primer extension were run in parallel with the sequence (lane G, A, T, and C) performed with the same oligonucleotide on U86. The sequence is reported on the left side: the arrow points to the XendoU cleavage sites. C, U86-containing precursor was incubated, in the presence of Mn⁺⁺ ions, in ONE (ONE), with XendoU produced by reticulocyte lysate (ret-XendoU/+Mn), or with reticulocyte lysate as such (lanes ret/+Mn). As control, pre-mRNA was incubated with XendoU produced by reticulocyte lysate in the absence of Mn⁺⁺ ions (ret-XendoU/-Mn). The numbers below indicate incubation times: 0 min (lane 1), 45 min (lanes 2).

EXPERIMENTAL PROCEDURES

Massive isolation of oocyte germinal vesicles and nuclear extract preparation

X. laevis germinal vesicles were isolated following the procedure by Gandini-Attardi et al. (Gandini-Attardi et al., 1990) and the nuclear extracts were prepared as already described (Caffarelli et al., 1994).

Purification of XendoU activity

XendoU was purified from oocyte nuclear extracts (ONE). ONE was fractionated by ammonium sulphate precipitation. Solid ammonium sulphate (280 mg/ml) was slowly added to the nuclear extract up to 45% saturation and the suspension was stirred for 30' at 4°C and then centrifuged at 12.000 rpm for 30' at 4°C. The supernatant was made 70% saturated by a further addition of ammonium sulphate (240 mg/ml). The suspension was stirred and centrifuged as above. The resulting pellet was dissolved in ONE buffer (25 mM Hepes pH 7.5, 50 mM NaCl, 0.1 mM EDTA, 1 mM DTT, 10% glycerol) and then applied onto an hydroxyapatite

column (CHT-II Econocolumn, Biorad). Column was washed with ONE buffer and then eluted with 5 column volumes of 100 mM Na-phosphate pH 7 in ONE buffer. The eluate was collected in 1 ml fractions next tested for the endonuclease activity. Selected fractions were pooled, diluted with 3 volumes of ONE buffer and applied on a Blue Sepharose column (Blue Sepharose Fast Flow Pharmacia). The column was washed with ONE buffer and then eluted with 5 column volumes of 0.2 M NaCl in ONE buffer. The eluate was then collected in 1 ml fractions; those displaying the specific activity were pooled and dialysed against ONE buffer. Protein mixture was subjected to a second fractionation on hydroxyapatite column. The elution was performed with 10 column volumes of a linear gradient 0-100 mM Na-phosphate pH 7 in ONE buffer. Collected fractions were tested and those with activity were pooled and concentrated by means of ultrafiltration device (Centricon C10, Millipore). The concentrated fractions were then applied on a gel-filtration column (Pharmacia) previously equilibrated in ONE buffer. Elution was monitored collecting 0.5 ml fractions which were tested for specific activity.

Considering the yield of purified protein it can be assumed that XendoU represents no more than 1/1000 of the protein mass present in nuclear extract. To obtain enough protein for sequencing and characterization the described procedure was carried out on several ONE samples of 15-20 ml collecting together the final purified fractions.

Preparation and isolation of tryptic peptides

Protein bands from SDS-PAGE analysis (5 µg) stained with Coomassie Blue R250 were excised, reduced with dithiothreitol and carboxamidomethylated. Gel pieces were equilibrated in 25 mM NH_4HCO_3 , pH 8 and finally digested *in situ* with trypsin at 37 °C for 18 h. Peptides were extracted by sonication with 100 µl of 25 mM NH_4HCO_3 /acetonitrile 1:1 v/v, pH 8 (twice). Peptide mixture was fractionated by reverse-phase HPLC on a Vydac C_{18} column 218TP52 (250 x 1 mm), 5 µm, 300 Å pore size (The Separation Group, USA) by using a linear gradient from 5% to 60% of acetonitrile in 0.1% TFA over 60 min, at flow rate of 90 µL/min. Individual components were manually collected and lyophilised.

Peptide Sequencing and Mass spectrometry analysis

Sequence analysis was performed using a Procise 491 protein sequencer (Applied Biosystems, USA) equipped with a 140C

microgradient apparatus and a 785A UV detector (Applied Biosystems, USA) for the automated identification of PTH-amino acids.

Matrix assisted laser desorption ionization mass spectra were recorded using a Voyager DE-PRO mass spectrometer (Applied Biosystems, USA); an analytical mixture containing α -cyano-4-hydroxycinnamic acid was applied to the sample plate and allowed to be dried. Mass calibration was performed using the molecular ions from peptides produced by tryptic auto-proteolysis and the matrix as internal standards.

In vitro RNA synthesis and oocyte microinjection

U16-containing precursor (003 RNA), including the third intron of the L4 r-protein gene of *X. laevis*, was transcribed from plasmid 003 digested with HindIII (Caffarelli et al., 1994). U86-containing precursor was obtained as already described (Filippini et al., 2001). *In vitro* transcription reactions were performed in the presence of (32 P) α UTP as described (Caffarelli et al., 1998) and pre-mRNAs were injected into nuclei of stage VI oocytes as already described (Caffarelli et al., 1994).

In vitro processing reactions

ONE assay: as described by (Caffarelli et al., 1994).

XendoU assay: the reaction mixture (25 μ l) contained 3×10^4 cpm of (32 P) labelled pre-mRNA, 6 mM MnCl_2 , 50 mM NaCl, 25 mM Hepes pH 7.5, 1 mM DTT, 10 μ g of E. Coli tRNA, 20 U of RNase inhibitor (PROMEGA) and 1 ng of purified XendoU. Reaction mixtures were incubated with RNA substrates at 24°C for indicated times. The products of the reactions were analysed on 6% polyacrylamide-7M urea gels.

Substrate specificity

The oligoribonucleotides

P1 (5'-GGAAACGUAUCCUUUGGGAG-3'), SEQ ID No 3;

P2 (5'-GGAAACGUAUCCUUGGGAGT-3'), SEQ ID No 4;

P3 (5'-GGAAACGUAUCCUCUGGGAG-3'), SEQ ID No 5;

P4 (5'-GGAAACGUAUCCUGUGGGAG-3'), SEQ ID No 6;

were 5' labelled: 10 pmol of each synthetic substrate were incubated at 37°C for 30 min, in the presence of 10 units of Polynucleotide Kinase (Roche), and 10 μ Ci of (32 P)Y-ATP. The reaction was terminated at 65°C for 5 min, primers were gel purified on 10% polyacrylamide-7M urea and incubated for 30 min in the presence of ONE or purified XendoU as described above. RNA was extracted and analysed on 10%

polyacrylamide-denaturing gel. RNA ladder was obtained by incubation of P4 oligo (200.000 cpm) in 500mM NaHCO₃ at 90°C for 20 min.

Analysis of 3' ends of cleavage products

³²P-labelled and gel purified I-1 molecules (I-1a and I-1b generated by cleavages at the major sites a and b upstream from U16, see scheme of Figure 1A), obtained *in vitro* after incubation of 003 RNA with ONE or XendoU, or *in vivo* after oocyte microinjection, were incubated in 10 µl of 10 mM HCl at 25°C for 2 hours to open the cyclic phosphate as described by Forster (Forster et al., 1990). The phosphate was then removed by incubation of the RNA in 50 mM Tris-HCl pH 8.5, 0,1 mM EDTA in the presence of 1 U of calf intestine Alkaline Phosphatase at 50°C for 60 min. The enzyme was inactivated by adding 1/10 volume of 0.2 M EGTA and the mixture heated at 65°C for 10 min. RNA was extracted with phenol/chloroform and analysed on 10% polyacrylamide-7M urea gel.

Isolation of XendoU cDNA

A *X. laevis* stage 28 embryo cDNA library, constructed in λZAP II vector, was screened using a specific probe obtained by PCR amplification on *X.laevis* cDNA with degenerated oligonucleotides (MAHs 5'-ATGGCICAYGAYTAYYTIGT-3', SEQ ID No 7 and IGTa 5'-ACIGGRTAIGCIGTICCIAT-3', SEQ ID No 8) designed on peptides obtained by tryptic digestion of purified XendoU.

XendoU cDNA expression in reticulocyte lysate

XendoU Open Reading Frame (ORF) was cloned into Blue Script vector and ³⁵(S)Methionine-labelled protein was produced by *in vitro* transcription and translation using the TnT-coupled Reticulocyte Lysate System Kit (PROMEGA) according to the manufacturer's instructions. Translational products were analysed on 10% SDS-PAGE.

Primer extension analysis

In vitro transcribed U86-containing precursor was obtained from a standard T7 reaction, but in the presence of 500 µM unlabelled UTP. The transcript was injected into *X. laevis* oocytes or incubated in ONE or with purified XendoU. The processing product I-4 RNA was gel-purified and reverse transcribed (SS pre-amplification system - GIBCO) with the 5' terminally labelled oligonucleotide UHindIII (5'-AAGCTTCTTCATGGCGGCTCGGCCAAT-3', SEQ ID No 9) complementary to 19 nucleotides at the 3' end of the downstream exon.

The elongated products were run in parallel with the sequence obtained with the same primer on U86-containing precursor.

RESULTS

Purification of XendoU from *X. laevis* oocyte nuclear extracts

5 The authors previously developed an *in vitro* system able to reproduce the release of U16 snoRNA from its host intron (Caffarelli et al., 1994). When ^{32}P -labelled U16-containing precursor is incubated in *X. laevis* oocyte nuclear extract, in the presence of Mn^{++} ions, specific products, originating from endonucleolytic cleavages, are found (Figure 1A and 1B): the I-2 and I-1 molecules derive from cleavage upstream from U16 coding region, while I-3 and I-4 molecules are produced by cleavage downstream from U16. When double cleavage occurs on the same pre-mRNA molecule, pre-U16 products accumulate, which are then converted to the mature snoRNA by exonucleotidic digestion

15 The procedure of biochemical characterization is indicated in Figure 1C, while the protein content of the fractions displaying XendoU activity is shown in Figure 1D. After several chromatographic steps, a single protein of 37 kDa molecular mass was obtained as showed in Figure 1D, lane 6. The elution of XendoU activity during the last purification step on gel filtration column is consistent with a monomeric protein of 37 kDa. The assay for testing the presence of XendoU activity was performed by incubating ^{32}P -labelled U16-containing precursor with aliquots of the different fractionation steps. Figure 1B shows the comparison of XendoU activity of unfractionated nuclear extracts (ONE) with that of the gel filtration column (XendoU). Since previously it was demonstrated the dependence of XendoU activity on Mn^{++} ions, this cofactor is always present in the reaction: in both cases the same primary cleavage products, I-2 and I-3 and their complementary molecules I-1 and I-4, are generated. Primer extension analysis performed on cleavage products I-2 and I-4 revealed that the purified enzyme cleaves intronic sequences at the same U-rich *in vivo* recognized areas (Caffarelli et al., 1994).

Characterization of XendoU cleavage

35 The analysis of the ion dependence of cleavage was carried out by incubating ^{32}P -labelled U16 containing precursor with ONE or with purified XendoU in the presence of different cations as shown in Figure 2. In both cases Cd^{++} , Zn^{++} , Ni^{++} , Co^{++} and Pb^{++} do not activate cleavage;

whereas Mn^{++} , and to a minor extent Mg^{++} , produce the appearance of the specific cleavage products. These results indicate a Mn^{++} ion requirement for XendoU full activity.

The substrate selectivity of XendoU was further addressed by incubating the purified enzyme with synthetic oligoribonucleotide (P1), containing the distal XendoU cleavage site (site d, Figure 1A), localised upstream from U16, and with mutated derivatives thereof (P2 and P3). The obtained results, shown in Figure 3A, indicate that XendoU displays the same selectivity observed *in vivo* and that the minimal consensus cleavage site is an uracyl dimer.

The incubation of the enzyme with double stranded oligoribonucleotides, of 21 nucleotides, containing U-rich sequences demonstrated that only single stranded RNAs are recognized by XendoU.

The chemistry of XendoU cleavage was then assessed by determining the chemical nature of the ends of the cleaved products. To this aim we analysed the ends of ^{32}P -labelled I-1a and I-1b molecules produced with ONE or with the purified XendoU: these molecules were gel purified and treated with alkaline phosphatase, HCl, or both. Figure 3B shows that a slight decrease in migration, due to the loss of a negative charge, is obtained when the alkaline phosphatase treatment follows the HCl treatment (lanes 3). This result is obtained both with ONE (lanes ONE) and purified XendoU (lanes XendoU) and indicates that the 3' end of the molecules bears a phosphate group which is present in a 2'-3' cyclic form (Lund and Dahiberg, 1992; Forster et al., 1990). In fact, only after the acidic treatment the phosphate group can be removed by phosphatase resulting in a slight electrophoresis delay. This effect is observed both for the I-1a and I-1b molecules obtained with unfractionated oocyte nuclear extract and with XendoU. As previously reported (Caffarelli et al., 1996), the products of primary cleavage such as I-1 molecules, are *in vivo* unstable being rapidly degraded, after cleavage. Nevertheless, after very short incubation times, little amounts of I-1b molecules can be purified and subjected to the same treatment as described above. Figure 3B (lanes *in vivo*) indicates that also in this case a slight delay in migration is obtained (lane 3), demonstrating that the products of the *in vivo* reaction display 2'-3' cyclic phosphates as well.

Isolation of cDNA for XendoU

After purification, protein samples from SDS-PAGE were reduced, alkylated and digested with trypsin as reported in the experimental section. The resulting peptide mixture was resolved by R-HPLC and selected peptide fractions were submitted to automated Edman degradation. The three sequence portions determined are reported in Figure 4 (indicated as 1, 2 and 3). From these amino acid sequences, degenerated oligonucleotides were derived and employed in different combinations and different orientation in PCR amplification reactions on cDNA from polyA⁺ RNA extracted from *X. laevis* oocytes. Only the reaction performed with sequence 1 (forward) and sequence 3 (reverse) resulted in an amplification product (500 bp). Sequencing of this product indicated the presence of an Open Reading Frame containing peptide 2. This cDNA probe was then utilised for the screening of a *X. laevis* stage 28 embryo cDNA library, allowing the isolation of a full-length cDNA (figure 4). The amino acid sequence determined was confirmed by MALDI-MS spectra of the tryptic peptides. In fact, signals observed at m/z 565.29, 814.45, 1004.48, 10025.54, 1132.59, 1190.60, 1490.78, 1504.80, 1520.70, 1729.91, 1758.82, 1988.08, 2000.00, 2014.01, 2076.99, 2162.98, 2234.14, 2238.05, 2394.15, 2432.26, 3058.51 and 3370.66 were ascribed to peptides 196-200, 126-132, 117-124, 6-14, 117-125, 41-52, 114-125, 260-271, 15-26, 137-149, 53-67, 256-271, 53-69, 275-292, 204-220, 150-169, 133-149, 171-188, 170-188, 117-136, 6-31 and Ac4-31. This result allowed to cover 65% of the entire sequence, explaining the reluctance of a blotted protein sample to Edman degradation.

Cloning and expression of XendoU cDNA

XendoU ORF, 876 bp, was cloned into Blue Script vector and the protein was produced by *in vitro* transcription and translation using reticulocyte lysate. The translational product was analysed on SDS-PAGE revealing a protein of 37 kDa molecular mass (Figure 5 A). In order to assess the nature of the 37 kDa protein, enzymatic activity was assayed by incubating the ³²P-labelled RNA substrate in reticulocyte lysate expressing XendoU ORF. The activity assay was carried out in parallel with the unfractionated extract: Figure 5B shows that the cleavage products generated by the translated 37 kDa protein (lane ret-XendoU/+Mn) exactly match those obtained with the extract (lanes ONE). Furthermore, the lack of cleavage when Mn⁺⁺ ions are not added to the

reaction mixture (lane ret-XendoU/-Mn) confirms the specific ion requirement for XendoU and suggests that the binding to the cofactor is reversible. As negative control the activity assay was carried out by incubating RNA substrate in reticulocyte lysate as such. in the presence of Mn⁺⁺ ions (lane ret/+Mn).

XendoU also participates to U86 snoRNA biosynthesis

The authors previously identified a snoRNA, named U86, encoded by an intron of Nop56 gene of *X. laevis*. It was also shown that U86 has homologs both in human, where it displays the same genomic organization, and in yeast where it is embedded in the ORF of Rib1 gene (Filippini et al., 2001). As U16 snoRNA, also U86 is contained in a very poorly spliceable intron and its biosynthesis appears to be alternative to that of the co-transcribed mRNA. The injection of ³²P-labelled U86-containing precursor into *X. laevis* oocytes generates the truncated products I-2 and I-3 and their 5' and 3' complementary molecules, 1-1 and 1-4, by means of upstream and downstream cleavages from U86 coding region (Figure 6A, lanes *in vivo*).

Processing of U86-containing precursor with purified XendoU (Figure 6A, lanes XendoU) or with the reticulocyte lysate expressing XendoU ORF (Figure 6C, lane ret-XendoU/+ Mn) demonstrates that the enzyme is responsible for the occurring cleavage downstream from U86 coding region. The effector of the cleavage upstream from U86, that produces I-2 and I-1 molecules, is not yet known and it is lost in oocyte nuclear extracts (Figure 6A, lanes ONE). The XendoU cleavage sites, downstream from U86, were determined by primer extension on I-4 molecules and correspond to two U-rich sequences (Figure 6B).

BIBLIOGRAPHY

Allmang, C., KufEL, J., Chanfreau, G., Mitchell, P., Petfaiski E. and Tollervey, D. (1999) Functions of the exosome m rRNA, snoRNA and snRNA synthesis. EMBO J. 18, 5399-5410.

Ambros.V. (2001) Dicing up RNAs. Science 293, 811-813

Bachellerie, J.P., Cavaille, J. and Qu, L.H. (2000) Nucleotide modifications of eukaryotic rRNAs: the world of small nucleolar RNA guides revisited. The Ribosome: structure, function, antibiotics, and cellular interactions. (2000 ASM Press, Washington, D.C.) pp. 191-203.

Bachmann, M., Messer, R., Trautmann, F. and Muller, W.E.G. (1984) 12S small nuclear ribonucleoprotein- associated acidic and

pyrimidine-specific endoribonuclease from calf thymus and L5178y cells. Biochim. Biophys. Acta 783, 89-99.

Bernstein, E., Caudy, A.A., Hammond, S.M. and Hannon, G.J. (2001) Role for a bidentate ribonuclease in the initiation step of RNA interference. Nature 409, 363-366.

Bujnicki, J.M., and Rychlewski, L. (2000) Prediction of a common fold for all four subunits of the yeast tRNA splicing endonuclease: implications for the evolution of the EndA/Sen family. FEBS Lett. 486, 328-9.

Caffarelli, E., Losito, M., Giorgi, C., Fatica, A., and Bozzoni, I. (1998) *in vivo* identification of nuclear factors interacting with the conserved elements of box C/D small nucleolar RNAs. Mol. Cell. Biol. 2, 1023-1028.

Caffarelli, E., Arese, M., Santoro, B., Fragapane, P., and Bozzoni, I. (1994) *In vitro* study of processing of the intron-encoded U16 small nucleolar RNA in *Xenopus laevis*. Mol. Cell. Biol. 14, 2966-2974

Caffarelli, E., Fatica, A., Prislei, S., De Gregorio, E., Fragapane, P., and Bozzoni, I. (1996) Processing of the intron-encoded U16 and U18 snoRNAs: the conserved c and D boxes control both the processing reactions and the stability of the mature snoRNA. EMBO J. 5, 1121-1131.

Castano, J.G., Tobian, J.A., and Zasloff, M. (1985) Purification and characterization of an endonuclease from *Xenopus laevis* ovaries which accurately processes the 3' terminus of human pre-tRNA-Met(i) (3' pre-tRNase). J. Biol. Chem. 260, 9002-9008

Chanfreau, G., Elela, S. A., Ares, M.Jr., and Guthrie, C. (1997) Alternative 3'-end processing of U5 snRNA by RNase III. Genes Dev. 11, 2741-2751

Chanfreau, G., Legrain, P., and Jacquier, A. (1998) Yeast RNase III as a key processing enzyme in small nucleolar RNAs metabolism. J Mol. Biol. 284, 975-88.

Cirino, N. M., Cameron, C.E., Smith, J.S., Rausch, J.W., Roth, M. J., Benkovic, S.J. and Le Grice, S.F. (1995) Divalent cation modulation of the ribonuclease functions of human immunodeficiency virus reverse transcriptase. Biochemistry 34, 9936-9943.

Court, D. (1993) RNase III: a double-strand processing enzyme. In: Brawerman, G. and Belasco, J., eds Control of mRNA stability. (New York, Academic, Press), pp.70-116.

Dange, V., Van Atta, R. B. and Hecht, S.M. (1990) A Mn²⁺-dependent ribozyme. *Science* 248, 585-588.

Deutscher, M. P. (1985) E.coli RNAses: making sense of alphabet soup. *Cell* 40, 731-732.

5 Deutscher, M. P. (1993) Ribonuclease multiplicity, diversity, and complexity, *J. Biol. Chem.* 268, 13011-13014

Eder, P.S. and Walder, J.A. (1991) Ribonuclease H from K562 human erythroleukemia cells. Purification, characterization, and substrate specificity. *J. Biol. Chem.* 266,6472-6479,

10 Elela, S.A., Igel, H., and Ares. M.Jr. (1996) Rnase III cleaves eukaryotic preribosomal RNA at a U3 snoRNP-dependent site. *Cell*, 85,115-124.

Filippini, D., Renzi, F., Bozzoni, I., and Caffarelli.E. (2001) U86, a novel snoRNA with an unprecedented gene organisation in yeast. *Biochem. Biophys. Res. Commun.* 288, 16-21

15 Forster,A.C., Davies, C., Hutchins, C.J. and Symons R.H. (1990) Characterization of self-cleavage of viroid and virusoid RNAs. *Methods Enzymol.* 181, 583-607

20 Fragapane, P., Prislei, S., Michienzi, A., Caffarelli, E. and Bozzoni I. (1993) A novel small nucleolar RNA (U16) is encoded inside a ribosomal protein intron and originates by processing of the pre-mRNA. *EMBO J.* 12, 2921-2928.

25 Freemont, P.S., Friedman, J.M., Beese, L.S., Sanderson M.R. and Steitz, T. A. (1988) Cocystal structure of an editing complex of Klenow fragment with DNA *Proc. Natl. Acad. Sci. USA* 85, 8924-8928.

Gandini-Attardi, D., Margarit, I. and Tocchini-Valentini, G.P. (1985) Structural alteration in mutant precursors of the yeast tRNA₃^{Leu} gene which behave as detective substrates for a highly purified splicing endoribonuclease. *EMBO J.* 4,3289-3297.

30 Gandini-Attardi, D., Baldi, I.M., Mattoccia, E., and Tocchini-Valentini G.P. (1990) Transfer RNA splicing endonuclease from *Xenopus laevis*. *Methods Enzymol.* 181, 510-517.

35 Giorgi, C., Fatica, A., Nagel, R., and Bozzoni, I. (2001) Release of U18 snoRNA from its host intron requires interaction of Nop1p with the Rnt1p endonuclease. *EMBO J.* 20, 6856-65.

Hirose, T., and Steitz J.A. (2001) Position within the host intron is critical for efficient processing of box C/D snoRNAs in mammalian cells. *Proc. Natl. Acad. Sci. USA* 98, 12914-12919.

5 Kazakov, S. and Altman, S. (1992) A trinucleotide can promote metal ion-dependent specific cleavage of RNA. *Proc. Natl. Acad. Sci. USA* 89, 7939-7943.

Kufel, J., Dichtl, B., and Tollervey, D. (1999) Yeast Rntip is required for cleavage of the pre-ribosomal RNA in the 3' ETS but not the 5' ETS. *RNA* 5, 909-917

10 Lygerou, Z., Allmang, C., Tollervey, D., and Seraphin, B. (1996) Accurate processing of a eukaryotic precursor ribosomal RNA by ribonuclease MRP *in vitro*. *Science* 272, 268-270

Lund, E. and Dahlberg J.E. (1992) Cyclic 2',3'-phosphates and nontemplated nucleotides at the 3' end of spliceosomal U6 small nuclear RNA's. *Science* 255, 327-330

Maxwell, E.S. and Fournier, M.J. (1995) The small nucleolar RNAs. *Annu. Rev. Biochem.* 64, 897-934

20 McDowall, K.J., Kaberdin, V.R., Wu, S.W., Cohen, S.N. and Lin-Chao, S. (1995) Site specific RNase E cleavage of oligonucleotides and inhibition by stem-loops. *Nature* 374, 287-290

Nashimoto, M., (1995) Conversion of mammalian tRNA 3' processing endoribonuclease to four-base-recognising RNA cutters. *Nucl. Acids Res.* 23, 3642-3647

25 Nicholson, A.W. (1997) in *Ribonucleases: structures and functions*, eds. D'Alessio, G. and Riordan, J.F. (Academic New York), pp. 1-49

Pan, T., Long, D. M. and Uhlenbeck, O.C. (1993) *The RNA World*, ed. R.F. Gesteland and J.F. Atkins, CSHL Press, 271-302.

30 Peebles, C. L., Gegenheimer, P., and Abelson, J. (1983) Precise excision of intervening sequences from precursor tRNAs by a membrane-associated yeast endonuclease. *Cell* 32, 525-536.

35 Prislei, S., Sperandio, S., Fragapane, P., Caffarelli, E., Presutti, C. and Bozzoni, I. (1992) The mechanisms controlling ribosomal protein L1 pre-mRNA splicing are maintained in evolution and rely on conserved intron sequences. *Nucl. Acids Res.* 17, 4473-4479

Prislei, S., Fatica, A., De Gregorio, E., Arese, M., Fragapane, P., Caffarelli, E., Presutti, C. and Bozzoni, I. (1995) Self-cleaving motifs

are found in close proximity to the sites utilised for U16 snoRNA processing. *Gene* 163,221-226.

- 5 Rauhut, R., Green, P.R., and Abelson, J. (1990). Yeast tRNA-splicing endonuclease is a heterotrimeric enzyme. *J. Biol. Chem.* 265,18180-18184.

- Schoenberg, D.R., and Chemokaiskaya, E. (1997) Ribonucleases involved in eukaryotic mRNA turnover, p. 217-240. In J. Harford and D. R. Morris (ed.), *mRNA metabolism and post-transcriptional gene regulation*. Wiley, New York; N. Y.

- 10 Steitz, J. A. and Steitz, T. A. (1993) A general two-metal-ion mechanism for catalytic RNA. *Proc. Natl. Acad. Sci. USA* 90, 6498-6502.

- Trotta, C.R., Miao, F., Am, E.A., Stevens, S.W., Ho, C.K., Rauhut, R., and Abelson, J.N. (1997) The yeast tRNA splicing endonuclease: a tetrameric enzyme with two active site subunits homologous to the archaeal tRNA endonucleases. *Cell* 89, 849-58

- 15 Zamore P.D. (2001) Thirty-three years later, a glimpse at the ribonuclease III active site. *Molecular Cell* 8,1158-1160

- Watkins, N.J., Segault, V., Charpentier, B., Nottrott, S., Fabrizio, P., Bachi, A., Wilm, M., Rosbash, M., Branlant, C., and Luhrmann, R. (2000) A common core RNP structure shared between the small nucleolar box C/D RNPs and the spliceosomal U4 snRNP. *Cell* 103, 457-466.

20

CLAIMS

1. Nucleic acid encoding for a protein with endoribonucleasic activity wherein said protein with endoribonucleasic activity is characterized in that it is polyU specific, Mn^{++} ions dependent and able to release 2'-3' cyclic phosphate and 5'OH RNA ends cleavage products.

5

2. Nucleic acid according to claim 1 substantially including nucleotide sequence of figure 4, functional homologs thereof or a complementary sequence thereto.

3. Recombinant vector able to express effectively the inventive nucleic acid in prokaryotes according to claims 1 or 2.

10

4. Recombinant vector able to express effectively the inventive nucleic acid in eukaryotes according to claims 1 or 2.

5. Protein with endoribonucleasic activity characterized in that it is polyU specific, Mn^{++} ions dependent, and able to release 2'-3' cyclic phosphate and 5'OH RNA ends cleavage products or functional portions thereof.

15

6. Protein according to claim 5 encoded by nucleic acid according to claims 1 or 2,

7. Protein according to claim 6 having substantially the amino acid sequence of figure 4.

20

PURIFICATION, CLONING AND BIOCHEMICAL CHARACTERIZATION
OF XendoU, ENDORIBONUCLEASIC ACTIVITY INVOLVED IN SMALL
NUCLEAR RNA SPLICING-INDEPENDENT BIOSYNTHESIS IN
XENOPUS LAEVIS

5

ABSTRACT

Cloning and characterization of the gene for Mn^{++} dependent
RNA-se endoribonucleasic activity able to generate 2'-3' cyclic phosphate
and 5'OH ends.

10

SEQUENCE LISTING

<110> BOZZONI, IRENE
CAFFARELLI, ELISA
LANEVE, PIETRO

<120> PURIFICATION, CLONING AND BIOCHEMICAL CHARACTERIZATION
OF XENDOU, ENDORIBONUCLEASIC ACTIVITY INVOLVED IN SMALL
NUCLEAR RNA SPLICING-INDEPENDENT BIOSYNTHESIS IN
XENOPUS LAEVIS

<130> 2520-1050

<140> 10/520,401

<141> 2005-09-12

<150> PCT/IT03/00424

<151> 2003-07-04

<150> IT RM2002A000365

<151> 2002-07-08

<160> 10

<170> PatentIn Ver. 3.3

<210> 1

<211> 1265

<212> DNA

<213> Xenopus laevis

<220>

<221> CDS

<222> (39)..(914)

<400> 1

attggggaac tgggagcaga gagtgacggg caggagcc atg gcg agt aac agg ggg 56
Met Ala Ser Asn Arg Gly
1 5

cag ctg aac cat gaa ctc tcc aag ctg ttt aat gag ctg tgg gac gca 104
Gln Leu Asn His Glu Leu Ser Lys Leu Phe Asn Glu Leu Trp Asp Ala
10 15 20

gat cag aac cgg atg aag tcc ggg aag gat tat cgg atc tcc ttg cag 152
Asp Gln Asn Arg Met Lys Ser Gly Lys Asp Tyr Arg Ile Ser Leu Gln
25 30 35

ggt aaa gca ggg tac gta ccc gcc ggt tcc aac cag gcc agg gac agc 200
Gly Lys Ala Gly Tyr Val Pro Ala Gly Ser Asn Gln Ala Arg Asp Ser
40 45 50

gcc tcg ttc ccg ctc ttc cag ttc gtc gat gag gag aag ctg aag agc 248
Ala Ser Phe Pro Leu Phe Gln Phe Val Asp Glu Glu Lys Leu Lys Ser
55 60 65 70

agg aag acg ttt gca acc ttc att tcc ctg ctg gac aat tat gag atg	296
Arg Lys Thr Phe Ala Thr Phe Ile Ser Leu Leu Asp Asn Tyr Glu Met	
75 80 85	
gac acg ggg gtg gcc gag gtt gtg act ccg gag gaa atc gct gaa aac	344
Asp Thr Gly Val Ala Glu Val Val Thr Pro Glu Glu Ile Ala Glu Asn	
90 95 100	
aac aac ttc ctg gac gcc att ctg gaa acc aaa gtg atg aag atg gca	392
Asn Asn Phe Leu Asp Ala Ile Leu Glu Thr Lys Val Met Lys Met Ala	
105 110 115	
cat gac tac ctg gtg agg aag aac caa gcc aaa ccc acc cgg aat gac	440
His Asp Tyr Leu Val Arg Lys Asn Gln Ala Lys Pro Thr Arg Asn Asp	
120 125 130	
ttc aag gtc caa ctg tac aac atc tgg ttc cag ctg tac tca cgg gcc	488
Phe Lys Val Gln Leu Tyr Asn Ile Trp Phe Gln Leu Tyr Ser Arg Ala	
135 140 145 150	
cca ggg agc aga ccc gat tgc tgc ggc ttt gag cac gtg ttt gtg gga	536
Pro Gly Ser Arg Pro Asp Ser Cys Gly Phe Glu His Val Phe Val Gly	
155 160 165	
gaa tcg aag cga ggg cag gag atg atg ggg ctt cac aac tgg gtc cag	584
Glu Ser Lys Arg Gly Gln Glu Met Met Gly Leu His Asn Trp Val Gln	
170 175 180	
ttt tac ctt cag gag aag agg aag aac atc gac tat aaa gga tac gtg	632
Phe Tyr Leu Gln Glu Lys Arg Lys Asn Ile Asp Tyr Lys Gly Tyr Val	
185 190 195	
gct cgg cag aac aag agt cgg ccg gat gaa gat gat cag gtg ttg aac	680
Ala Arg Gln Asn Lys Ser Arg Pro Asp Glu Asp Asp Gln Val Leu Asn	
200 205 210	
ctg cag ttc aat tgg aag gag atg gtg aaa ccc gtc ggc agc agc ttc	728
Leu Gln Phe Asn Trp Lys Glu Met Val Lys Pro Val Gly Ser Ser Phe	
215 220 225 230	
att ggc gtc agc ccg gaa ttc gaa ttc gcc ctt tac acc atc gtc ttc	776
Ile Gly Val Ser Pro Glu Phe Glu Phe Ala Leu Tyr Thr Ile Val Phe	
235 240 245	
ctc gcg tct cag gag aag atg agc cga gaa gtc gtt cgg ctg gaa gaa	824
Leu Ala Ser Gln Glu Lys Met Ser Arg Glu Val Val Arg Leu Glu Glu	
250 255 260	
tac gaa ctg cag atc gtc gtc aat cgc cac ggc cgt tat ata ggg acc	872
Tyr Glu Leu Gln Ile Val Val Asn Arg His Gly Arg Tyr Ile Gly Thr	
265 270 275	
gcc tac ccc gtc ctc ctg agc acc aat aac ccg gat ctg tac	914
Ala Tyr Pro Val Leu Leu Ser Thr Asn Asn Pro Asp Leu Tyr	
280 285 290	
tgagggggcg gggctagaga tcacagccgg ttcccacggt ttgggtgcat ttactaacia	974

aactgcacca atgcaacaat gcaagcagat aatgggggca ggtccatata cctctgcttt 1034
 ccctagcgtg tgtggggcac attaacccta taactgtcac tcaactgcacc agaccatta 1094
 ttttaacccca caagggacat caagccagtg ccttggttatg agagagcgca gccggggcctt 1154
 ctctactgtg aaacttctgt attgtataga gtttacttgg tttcttcctc cagacaattt 1214
 cacttttttt ttgctttgcc ttttaaccatt aaaagtccat gacatttctg t 1265

<210> 2

<211> 292

<212> PRT

<213> *Xenopus laevis*

<400> 2

Met	Ala	Ser	Asn	Arg	Gly	Gln	Leu	Asn	His	Glu	Leu	Ser	Lys	Leu	Phe	1	5	10	15
Asn	Glu	Leu	Trp	Asp	Ala	Asp	Gln	Asn	Arg	Met	Lys	Ser	Gly	Lys	Asp	20	25	30	
Tyr	Arg	Ile	Ser	Leu	Gln	Gly	Lys	Ala	Gly	Tyr	Val	Pro	Ala	Gly	Ser	35	40	45	
Asn	Gln	Ala	Arg	Asp	Ser	Ala	Ser	Phe	Pro	Leu	Phe	Gln	Phe	Val	Asp	50	55	60	
Glu	Glu	Lys	Leu	Lys	Ser	Arg	Lys	Thr	Phe	Ala	Thr	Phe	Ile	Ser	Leu	65	70	75	80
Leu	Asp	Asn	Tyr	Glu	Met	Asp	Thr	Gly	Val	Ala	Glu	Val	Val	Thr	Pro	85	90	95	
Glu	Glu	Ile	Ala	Glu	Asn	Asn	Asn	Phe	Leu	Asp	Ala	Ile	Leu	Glu	Thr	100	105	110	
Lys	Val	Met	Lys	Met	Ala	His	Asp	Tyr	Leu	Val	Arg	Lys	Asn	Gln	Ala	115	120	125	
Lys	Pro	Thr	Arg	Asn	Asp	Phe	Lys	Val	Gln	Leu	Tyr	Asn	Ile	Trp	Phe	130	135	140	
Gln	Leu	Tyr	Ser	Arg	Ala	Pro	Gly	Ser	Arg	Pro	Asp	Ser	Cys	Gly	Phe	145	150	155	160
Glu	His	Val	Phe	Val	Gly	Glu	Ser	Lys	Arg	Gly	Gln	Glu	Met	Met	Gly	165	170	175	
Leu	His	Asn	Trp	Val	Gln	Phe	Tyr	Leu	Gln	Glu	Lys	Arg	Lys	Asn	Ile	180	185	190	
Asp	Tyr	Lys	Gly	Tyr	Val	Ala	Arg	Gln	Asn	Lys	Ser	Arg	Pro	Asp	Glu	195	200	205	
Asp	Asp	Gln	Val	Leu	Asn	Leu	Gln	Phe	Asn	Trp	Lys	Glu	Met	Val	Lys	210	215	220	

Pro Val Gly Ser Ser Phe Ile Gly Val Ser Pro Glu Phe Glu Phe Ala
 225 230 235 240
 Leu Tyr Thr Ile Val Phe Leu Ala Ser Gln Glu Lys Met Ser Arg Glu
 245 250 255
 Val Val Arg Leu Glu Glu Tyr Glu Leu Gln Ile Val Val Asn Arg His
 260 265 270
 Gly Arg Tyr Ile Gly Thr Ala Tyr Pro Val Leu Leu Ser Thr Asn Asn
 275 280 285
 Pro Asp Leu Tyr
 290

<210> 3
 <211> 20
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 3
 ggaaacguau ccuuugggag 20

<210> 4
 <211> 20
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 4
 ggaaacguau ccuugggagg 20

<210> 5
 <211> 20
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 5
 ggaaacguau ccucugggag 20

<210> 6
 <211> 20
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 6
 ggaaacguau ccugugggag

20

<210> 7
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<220>
 <221> modified_base
 <222> (6)
 <223> inosine

<220>
 <221> modified_base
 <222> (18)
 <223> inosine

<400> 7
 atggcncayg aytayytngt

20

<210> 8
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<220>
 <221> modified_base
 <222> (3)
 <223> inosine

<220>
 <221> modified_base
 <222> (9)
 <223> inosine

<220>
 <221> modified_base
 <222> (12)
 <223> inosine

<220>
 <221> modified_base
 <222> (15)
 <223> inosine

<220>
 <221> modified_base
 <222> (18)
 <223> inosine

<400> 8
 acnggrtang cngtnccnat

20

<210> 9
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 9
 aagctttcttc atggcggctc ggccaat

27

<210> 10
 <211> 15
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 10
 tcttttcatt cattt

15